

A Brief Review on Obesity and It's Negative Effects on Health

Nidhi Sharma

Assistant Professor, IEC College of Engineering & technology, Greater Noida

Abstract - Obesity is a very complex disease involving an excessive amount of body fat. It has become a serious worldwide problem now. More than 300 million adults are classified as obese and the global number is predicted to reach 700 million by 2015. In past, obesity was not a common condition among the population of China, however. Obesity and diabetes have become a growing challenge, and one million new diabetic cases are reported in China every year. Increasing evidence suggests that obesity is a risk factor for diabetes and chronic kidney diseases. As a marker of obesity, high body mass index (BMI) has been reported to be related with diabetic nephropathy (DN) and end-stage renal disease (ESRD).

Index Terms - Obesity, Heart disease, Diabetes, inheritance, Sleep apnoea, osteoarthritis.

INTRODUCTION

Obesity is a very complex disease involving an excessive amount of body fat. Obesity is not just a cosmetic concern. It is a medical problem that increases risk of other diseases and health problems, such as,

- Heart disease.
- Diabetes.
- High blood pressure.
- Certain cancers.

It is also very difficult to avoid obesity because it results from a combination of inherited factors, combined with the environment and personal diet and exercise choices. [1]

On the other hand, modest weight loss can improve or prevent the health problems which are associated with obesity. Dietary changes increased physical activity and behaviour changes also can helps to lose weight. [2]

It has become a serious worldwide problem in few decades. More than 300 million adults are classified as obese and the global number is predicted to reach 700

million by 2015. In the past, obesity was not a common condition among the population of China; however, this is no longer the case [3]. Obesity and diabetes have become a growing challenge, and one million new diabetic cases are reported in China every year. Increasing evidence suggests that obesity is a risk factor for diabetes and chronic kidney diseases. As a marker of obesity, high body mass index (BMI) has been reported to be related with diabetic nephropathy (DN) and end-stage renal disease (ESRD) [4].

SYMPTOMS

Obesity generally diagnosed when body mass index (BMI) is 30 or higher. For determination of body mass index, divide person's body weight in pounds by person's height in inches squared and multiply by 703. Or divide person's body weight in kilograms by person's height in meters squared.

BMI	WEIGHT STATUS
Below 18.5	Underweight
18.5-24.9	Normal
25.0-29.9	Overweight
30.0 and higher	Obesity

Body mass index (BMI) provides a reasonable estimate of body fat. [5]

CAUSES

Although there are genetic, behavioural, metabolic and hormonal influences on body weight, But Obesity occurs when a person takes more calories than they burn through exercise and normal daily activities. Human body stores these excess calories as fat. [6] Most Americans people's diets are very high in calories often from fast food and high-calorie beverages. People with obesity might eat more

calories before feeling full, feel hungry sooner, or eat more due to stress or anxiety.

1. Family Inheritance & Influences-

The genes you inherit from your parents may affect the amount of body fat you store, and where that fat is distributed. Genetics may also play a role in how efficiently your body converts food into energy, how your body regulates your appetite and how your body burns calories during exercise.

Obesity tends to run in families. That's not just because of the genes they share. Family members also tend to share similar eating and activity habits. [7]

2. Lifestyle Choices-

Unhealthy diet. A diet that's high in calories, lacking in fruits and vegetables, full of fast food, and laden with high-calorie beverages and oversized portions contributes to weight gain.

Liquid calories. People can drink many calories without feeling full, especially calories from alcohol. Other high-calorie beverages, such as sugared soft drinks, can contribute to significant weight gain.

Inactivity. If you have a sedentary lifestyle, you can easily take in more calories every day than you burn through exercise and routine daily activities. Looking at computer, tablet and phone screens is a sedentary activity. The number of hours you spend in front of a screen is highly associated with weight gain. [8]

3. Certain disease & medication-

In some people, obesity can be traced to a medical cause, such as Prader-Willi syndrome, Cushing syndrome and other conditions. Medical problems, such as arthritis, also can lead to decreased activity, which may result in weight gain.

Some medications can lead to weight gain if you don't compensate through diet or activity. These medications include some antidepressants, anti-seizure medications, diabetes medications, antipsychotic medications, steroids and beta blockers. [9]

4. Social & Economic issues-

Social and economic factors are linked to obesity. Avoiding obesity is difficult if you don't have safe areas to walk or exercise. Similarly, you may not have been taught healthy ways of cooking, or you may not have access to healthier foods. In addition, the people you spend time with may influence your weight —

you're more likely to develop obesity if you have friends or relatives with obesity. [10]

5. Age-

Obesity can occur at any age, even in young children. But as you age, hormonal changes and a less active lifestyle increase your risk of obesity. In addition, the amount of muscle in your body tends to decrease with age. Generally, lower muscle mass leads to a decrease in metabolism. These changes also reduce calorie needs and can make it harder to keep off excess weight. If you don't consciously control what you eat and become more physically active as you age, you'll likely gain weight. [11]

6. Other factors-

- Pregnancy. Weight gain is common during pregnancy. Some women find this weight difficult to lose after the baby is born. This weight gain may contribute to the development of obesity in women. Breast-feeding may be the best option to lose the weight gained during pregnancy.
- Quitting smoking. Quitting smoking is often associated with weight gain. And for some, it can lead to enough weight gain to qualify as obesity. Often, this happens as people use food to cope with smoking withdrawal. In the long run, however, quitting smoking is still a greater benefit to your health than is continuing to smoke. Your doctor can help you prevent weight gain after quitting smoking.
- Lack of sleep. Not getting enough sleep or getting too much sleep can cause changes in hormones that increase your appetite. You may also crave foods high in calories and carbohydrates, which can contribute to weight gain.
- Stress. Many external factors that affect your mood and well-being may contribute to obesity. People often seek more high-calorie food when experiencing stressful situations.
- Microbiome. Your gut bacteria are affected by what you eat and may contribute to weight gain or difficulty losing weight.
- Previous attempts to lose weight. Previous attempts of weight loss followed by rapid weight regain may contribute to further weight gain. This phenomenon, sometimes called yo-yo dieting, can slow your metabolism.

Even if you have one or more of these risk factors, it doesn't mean that you're destined to develop obesity. You can counteract most risk factors through diet, physical activity and exercise, and behaviour changes. [13]

COMPLICATIONS

People with obesity are more likely to develop a number of potentially serious health problems, including:

- Heart disease and strokes. Obesity makes you more likely to have high blood pressure and abnormal cholesterol levels, which are risk factors for heart disease and strokes.
- Type 2 diabetes. Obesity can affect the way your body uses insulin to control blood sugar levels. This raises your risk of insulin resistance and diabetes.
- Certain cancers. Obesity may increase your risk of cancer of the uterus, cervix, endometrium, ovary, breast, colon, rectum, esophagus, liver, gallbladder, pancreas, kidney and prostate.
- Digestive problems. Obesity increases the likelihood that you'll develop heartburn, gallbladder disease and liver problems.
- Gynecological and sexual problems. Obesity may cause infertility and irregular periods in women. Obesity also can cause erectile dysfunction in men.
- Sleep apnea. People with obesity are more likely to have sleep apnea, a potentially serious disorder in which breathing repeatedly stops and starts during sleep.
- Osteoarthritis. Obesity increases the stress placed on weight-bearing joints, in addition to promoting inflammation within the body. These factors may lead to complications such as osteoarthritis.
- Severe COVID-19 symptoms. Obesity increases the risk of developing severe symptoms if you become infected with the virus that causes coronavirus disease 2019 (COVID-19). People who have severe cases of COVID-19 may require treatment in intensive care units or even mechanical assistance to breathe. [15]

QUALITY OF LIFE

Obesity can diminish your overall quality of life. You may not be able to do things you used to do, such as

participating in enjoyable activities. You may avoid public places. People with obesity may even encounter discrimination.

Other weight-related issues that may affect your quality of life include:

- Depression
- Disability
- Sexual problems
- Shame and guilt
- Social isolation
- Lower work achievement [16]

PREVENTION

Whether you're at risk of obesity, currently overweight or at a healthy weight, you can take steps to prevent unhealthy weight gain and related health problems. Not surprisingly, the steps to prevent weight gain are the same as the steps to lose weight: daily exercise, a healthy diet, and a long-term commitment to watch what you eat and drink.[17]

- Exercise regularly. You need to get 150 to 300 minutes of moderate-intensity activity a week to prevent weight gain. Moderately intense physical activities include fast walking and swimming.
- Follow a healthy-eating plan. Focus on low-calorie, nutrient-dense foods, such as fruits, vegetables and whole grains. Avoid saturated fat and limit sweets and alcohol. Eat three regular meals a day with limited snacking. You can still enjoy small amounts of high-fat, high-calorie foods as an infrequent treat. Just be sure to choose foods that promote a healthy weight and good health most of the time.
- Know and avoid the food traps that cause you to eat. Identify situations that trigger out-of-control eating. Try keeping a journal and write down what you eat, how much you eat, when you eat, how you're feeling and how hungry you are. After a while, you should see patterns emerge. You can plan ahead and develop strategies for handling these types of situations and stay in control of your eating behaviors.
- Monitor your weight regularly. People who weigh themselves at least once a week are more successful in keeping off excess pounds. Monitoring your weight can tell you whether your

efforts are working and can help you detect small weight gains before they become big problems.

- Be consistent. Sticking to your healthy-weight plan during the week, on the weekends, and amidst vacation and holidays as much as possible increases your chances of long-term success.[18]

REFERENCE

- [1] Ji CY, Cheng TO: Epidemic increase in overweight and obesity in Chinese children from 1985 to 2005. *Int J Cardiol* 2009, 132(1):1–10.
- [2] Hsu CY, McCulloch CE, Iribarren C, Darbinian J, Go AS: Body mass index and risk for end-stage renal disease. *Ann Intern Med* 2006, 144(1):21–28.
- [3] Hsieh MC, Hsieh YT, Cho TJ, Chen JF, Lin SD, Chen HC, Tu ST: Remission of diabetic nephropathy in type 2 diabetic Asian population: role of tight glucose and blood pressure control. *Eur J Clin Invest* 2011, 41(8):870–878.
- [4] Luk A, Chan JC: Diabetic nephropathy—what are the unmet needs? *Diabetes Res Clin Pract* 2008, 82(Suppl 1):S15–S20.
- [5] Speakman JR, Westerterp KR: Reverse epidemiology, obesity and mortality in chronic kidney disease: modelling mortality expectations using energetics. *Blood Purif* 2010, 29(2):150–157.
- [6] Valocikova I, Valocik G, Kristofova B, Druzbacka L: Obesity paradox and chronic kidney disease. *Bratisl Lek Listy* 2011, 112(7):402–406.
- [7] Pei YP, Greenwood CM, Chery AL, Wu GG: Racial differences in survival of patients on dialysis. *Kidney Int* 2000, 58(3):1293–1299.
- [8] Zhou BF: Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults—study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci* 2002, 15(1):83–96.
- [9] Li LS, Liu ZH: Epidemiologic data of renal diseases from a single unit in China: analysis based on 13,519 renal biopsies. *Kidney Int* 2004, 66(3):920–923.
- [10] Alberti KG, Zimmet PZ: Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998, 15(7):539–553.
- [11] Jennette JLO JC, Schwartz MM, Silva FG: *Heptinstall's Pathology of the Kidney*. 5th edition. Lippincott Williams & Wilkins; 1998.
- [12] Huang F, Yang Q, Chen L, Tang S, Liu W, Yu X: Renal pathological change in patients with type 2 diabetes is not always diabetic nephropathy: a report of 52 cases. *Clin Nephrol* 2007, 67(5):293–297.
- [13] Tone A, Shikata K, Matsuda M, Usui H, Okada S, Ogawa D, Wada J, Makino H: Clinical features of non-diabetic renal diseases in patients with type 2 diabetes. *Diabetes Res Clin Pract* 2005, 69(3):237–242.
- [14] Lemann J, Bidani AK, Bain RP, Lewis EJ, Rohde RD: Use of the serum creatinine to estimate glomerular filtration rate in health and early diabetic nephropathy. Collaborative study group of angiotensin converting enzyme inhibition in diabetic nephropathy. *Am J Kidney Dis* 1990, 16(3):236–243.
- [15] Chen HM, Chen Y, Zhang YD, Zhang PP, Chen HP, Wang QW, Li LS, Liu ZH: Evaluation of metabolic risk marker in obesity-related glomerulopathy. *J Ren Nutr* 2011, 21(4):309–315.
- [16] Lane PH, Steffes MW, Mauer SM: Estimation of glomerular volume: a comparison of four methods. *Kidney Int* 1992, 41(4):1085–1089.
- [17] Chen HM, Liu ZH, Zeng CH, Li SJ, Wang QW, Li LS: Podocyte lesions in patients with obesity-related glomerulopathy. *Am J Kidney Dis* 2006, 48(5):772–779.
- [18] Gellman DD, Pirani CL, Soothill JF, Muehrcke RC, Kark RM: Diabetic nephropathy: a clinical and pathologic study based on renal biopsies. *Medicine (Baltimore)* 1959, 38:321–367.